

**TUBERCULOSIS MEDICATION ORDER FOR DIRECTLY OBSERVED THERAPY (DOT)**

DOT is the observation of a person taking his/her prescribed medication for the treatment of tuberculosis (TB) disease or TB infection. Public Health guidelines recommend DOT as standard treatment for TB disease. *DOT can be done in-person or with the use of mobile video technologies.*

Client's Name: \_\_\_\_\_ D.O.B: \_\_\_\_\_

Medical Record No: \_\_\_\_\_ Allergies: \_\_\_\_\_

Public Health Nurse: \_\_\_\_\_ Telephone: \_\_\_\_\_

Date for DOT to begin: \_\_\_\_\_ Weight (kg) : \_\_\_\_\_

Medication	Strength (mg)	Number of Tabs	Total Dosage	Route	Frequency	Date Started	Date Stopped
Isoniazid							
Rifampin							
Ethambutol							
Pyrazinamide							
Vitamin B6							

If treating **PANSENSITIVE** Tuberculosis:

- Discontinue Ethambutol when DST result received
- Discontinue Pyrazinamide after 8 weeks dosing and/or 40 DOT doses

(Note: Public Health Nurse will confirm with Provider when each drug is being discontinued)

Comments: \_\_\_\_\_

Provider Signature: \_\_\_\_\_ Date: \_\_\_\_\_

Print Name: \_\_\_\_\_ Facility: \_\_\_\_\_

Phone No: \_\_\_\_\_ Fax No: \_\_\_\_\_